

Xylazine Wound Care Among PWUD Accessing Massachusetts Syringe Service Programs

¹Samia A Ismail BS, ²Margaret Shang MD, ³Stephen Murray MPH, NRP, ⁴Cristina Murray-Krezan PhD, ⁴Yihao Zheng MS, ⁴Gary McMurtrie BA, ^{2,4}Jane Liebschutz MD, MPH, ^{2,4}Raagini Jawa MD, MPH, FASAM

¹Boston University School of Public Health, Boston, MA, ²University of Pittsburgh Medical Center, Pittsburgh, PA, ³Boston Medical Center, Boston, MA, ⁴Center for Research in Healthcare, Division of General Internal Medicine, University of Pittsburgh, Pittsburgh, PA

Background

- Xylazine is an adulterant increasingly prevalent in the illicit opioid supply in Massachusetts (MA)
- People who use drugs (PWUD) exposed to xylazine often present with distinct wounds that can heal with medical care and self-treatment
- Understanding medical wound care experiences and wound self-treatment practices is critical to preventing further harm

Methods



175 PWUD past-year wounds at 3 MA syringe service programs (SSPs) in August 2023

33-item anonymous Qualtrics survey of multiple choice and free text questions on:

- ✓ Drug use practices
- ✓ Xylazine exposure
- ✓ Past experiences seeking medical care for wounds
- ✓ Past 90-day wound development
- ✓ Wound self-treatment practices

Responses analyzed using descriptive statistics and inductive content analysis

Participants given \$25 compensation



Outcomes Assessed

- PWUD experiences receiving medical care for wounds
- PWUD wound self-treatment practices


Results

1

Medical Wound Care Experiences

- N=171, 63% male, 63% white, 67% non-Hispanic
- 87% (N=148) had a suspected past 90-day xylazine wound
- Majority reported provider stigma, inadequate withdrawal management, and long wait times

Figure 1. Barriers to medical wound care experienced by PWUD with xylazine wounds (N=148)



“Nurses accuse me of lying about not injecting because I have a wound.”

2

Wound Self-Treatment Practices

Figure 2. Wound cleansers used by PWUD with xylazine wounds (N=148)





Figure 3. Wound coverage techniques used by PWUD with xylazine wounds (N=148)



“I cover my wound with whatever I have at work toilet paper, duct tape, paper towels.”

Conclusion

- Majority of PWUD seeking services at MA SSPs reported experiencing xylazine wounds concerning for xylazine exposure
- Interrupted or delayed medical care may contribute to advanced wound progression
- PWUD with xylazine wounds face several barriers to receiving medical care
- Use of caustic cleansing agents by PWUD with xylazine wounds was common and could slow healing

Key Takeaways

- Decreasing stigma and managing withdrawal symptoms makes medical care accessible at earlier stages of wound development
- Standardized xylazine wound self-care information is urgently needed

References

- Friedman J, Montero F, Bourgois P, et al. Xylazine spreads across the US: A growing component of the increasingly synthetic and polysubstance overdose crisis. *Drug Alcohol Depend.* 2022;233:109380. doi:10.1016/j.drugalcdep.2022.109380
- Ozga JE, Syvertsen JL, Zweifler JA, Pollini RA. A community-based study of abscess self-treatment and barriers to medical care among people who inject drugs in the United States. *Health Soc Care Community.* 2022;30(5):1798-1808. doi:10.1111/hsc.13559
- Ruiz-Colón K, Chavez-Arias C, Díaz-Alcalá JE, Martínez MA. Xylazine intoxication in humans and its importance as an emerging adulterant in abused drugs: A comprehensive review of the literature. *Forensic Sci Int.* 2014;240:1-8. doi:10.1016/j.forsciint.2014.03.015
- Torruebla RA. Xylazine (veterinary sedative) use in Puerto Rico. *Subst Abuse Treat Prev Policy.* 2011;6:7. Published 2011 Apr 11. doi:10.1186/1747-597X-6-7

Disclosures: None

Supported by: NIDA K12DA050607 (RJ, JML), 2023 Boston University School of Public Health Impact Grant (SI)

Acknowledgement: Our sincere gratitude to the persons who use drugs and front-line harm reduction organizations for their generosity in assisting with this research.